



Relating subsidence to land-use history using remote-sensing derived data for the Mekong delta, Vietnam

Philip S.J. Minderhoud^{1,2,*}, Laura Coumou^{1,#}, L.E. Erban^{3,4}, H. Middelkoop¹, E. Stouthamer¹, E.A. Addink¹

¹ Department of Physical Geography, Utrecht University, The Netherlands

² Department of Subsurface and Groundwater Systems, Deltares Research Institute, Utrecht, The Netherlands

³ Department of Earth System Science, Stanford University, Stanford

⁴ US EPA Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, Narragansett, RI, USA

* Corresponding author: P.S.J.Minderhoud@uu.nl



Presenting author

Introduction


The Vietnamese Mekong delta is subsiding due to a combination of natural and human-induced causes. Over the past several decades, large-scale anthropogenic land-use changes have taken place as a result of increased agricultural production, population growth and urbanization in the delta. Land-use changes can alter the hydrological system or increase loading of

the delta surface, amplifying natural subsidence processes or creating new anthropogenic subsidence. We quantified subsidence rates for the various land-use classes and past land-use changes and evaluated the relationship strength between current land use, land-use history and subsidence by predicting subsidence rates during the measurement period solely based on land-use history.

Data and approach

We created a new consistent time series of land-use maps by classifying Landsat Thematic Mapper (TM) 5 images using object-based image analysis. The land-use maps were used to create the land-use history for the period 1988-2006. Combined with InSAR-based subsidence rate, the subsidence rate for each individual land use and land-use change trajectory was quantified. To assess the strength of the relationship between land use, land-use change and subsidence, we evaluated the ability to predict subsidence rates based solely on land-use history.

Figure 1. Study area with InSAR-derived subsidence rates (2006-2010 by Erban et al., 2014) in the Vietnamese Mekong delta (VMD) in Southeast Asia.



Land-use maps




Figure 2. Land-use maps of the northeastern part of the Vietnamese Mekong delta derived from Landsat 5 TM imagery of 1988, 1996, 2006 and 2009.

Subsidence rate per land-use type




Figure 3. Estimated impact of subsidence drivers, and mean InSAR-based subsidence rate per land-use type (similar land-use class from 1988-2006). The estimated impact of each subsidence driver is ranked: minimal (-), low (\pm), moderate (+) and high (++) .

The effect of land-use change on subsidence rate

	Land use change		Subsidence rate (mm yr ⁻¹)			
	Original land use	Land use after change	Transition from LU 1 to LU 2			
	Land use 1	Land use 2	Unchanged LU 1	1996-2006	1988-1996	Unchanged LU 2
Development	Marshland	Dry-season rice	6	8	8	8
	Cultivating	Marshland	6	8	8	9
	undeveloped	Marshland	6	9	10	13
	land	Marshland	6	9	11	14
Changing	Mangroves	Aquaculture	13	11	11	11
	agriculture	Dry-season bare	13	10	8	8
		Dry-season rice	8	10	12	14
		Orchard	8	9	27	20
Urbanization	Dry-season bare	Urban dense	13	16	23	20
	Orchard	Urban dense	14	16	20	20

Figure 4. Impact of past land-use (LU) changes on the subsidence rates in the Mekong delta. The mean subsidence rates (in mm yr⁻¹) for areas in which LU 1 and LU 2 was unchanged during the period 1988-2006 (columns 'Unchanged LU 1' and 'Unchanged LU 2') and for areas that experienced a transition from LU 1 to LU 2 between 1996-2006 (column 'Transition from LU 1 to LU 2 1996-2006') and 1988-1996 (column 'Transition from LU 1 to LU 2 1988-1996'), respectively <10 and 10-18 years before the measurement period (2006-2010).

Land-use change

Main land-use classes	Land-use maps		
	1988	1996	2006
Undeveloped	18	9	5
Rice	54	50	54
Other agriculture	21	32	23
Aquaculture	4	4	8
Urban	3	5	10

Table 1. Area in percentages of the main land-use classes in the land-use maps of 1988, 1996 and 2006. Color coded from small (green) to large (red).

Subsidence prediction




Figure 5. A) Observed InSAR-based subsidence rate, B) predicted subsidence rates for the period 2006-2010, and C) difference between observed and predicted subsidence. The predicted subsidence over the period 2006-2010 (B) is based on land-use history. 66% to 92% of the predictions fall the error range of the InSAR-based subsidence (5 to 10 mm) of the observed values, observations.

Conclusions

- Land use and land-use history have an indirect causal relationship with subsidence rates in the Mekong delta
- Different land-use classes are experiencing different rates of subsidence
- Highest subsidence rates were found for land-use classes in which the natural environmental conditions were most altered by human activities

Acknowledgements

The Rise and Fall project is funded by NWO-WOTRO (W 07.69.105), Deltares and TNO-Geological Survey of the Netherlands.

References

Erban, L. E., Gorelick, S. M., & Zebker, H. A. (2014). Groundwater extraction, land subsidence, and sea-level rise in the Mekong Delta, Vietnam. *Environmental Research Letters*, 9(8), 1-6.